

Effect of postoperative kidney events on ICU and hospital length of stay

Outcome: ICU length of stay

- **aki\_0\_no\_1\_yes:**

factor	AME	lower	upper	p
aki_0_no_1_yes1	0.5895	- 0.5006	1.68	0.2892

- **Criterion\_aki\_1\_scr\_2\_diuresis\_3\_both:**

factor	AME	lower	upper	p
criteria_per_aki_1_scr_2_diuresis_3_both2	- 0.16 26	- 1.37	1.0 45	0.791 8
criteria_per_aki_1_scr_2_diuresis_3_both3	3.21 1	- 0.59 17	7.0 15	0.097 92

- **aki\_stage:**

factor	AME	lower	upper	p
aki_stage2	0.4184	- 0.7419	1.579	0.4797
aki_stage3	12.05	-3.619	27.72	0.1317

- **persistent\_aki\_0\_transient\_aki\_1\_persistent\_aki:**

factor	AME	lower	upper	p
persistent_aki_0_transient_aki_1_persistent_aki1	3.2 55	- 0.17 37	6.6 84	0.06 279

- **persistent\_aki\_stage:**

factor	AME	lower	upper	p
persistent_aki_stage1	1.278	- 0.2937	2.849	0.111
persistent_aki_stage2	5.5	-6.703	17.7	0.377
persistent_aki_stage3	13.5	-2.913	29.91	0.1069

- **akd\_0\_no\_1\_yes:**

factor	AME	lower	upper	p
akd_0_no_1_yes1	4.748	- 2.749	12.25	0.2145

Outcome: Hospital length of stay

- **aki\_0\_no\_1\_yes:**

factor	AME	lower	upper	p
aki_0_no_1_yes1	- 0.9414	- 4.018	2.135	0.5486

- **criterion\_aki\_1\_scr\_2\_diuresis\_3\_both:**

factor	AME	low er	upp er	p
crieterio_per_aki_1_scr_2_diuresis_3_both2	0.6105	-3.504	4.725	0.7712
crieterio_per_aki_1_scr_2_diuresis_3_both3	3.198	-3.58	9.975	0.3551

- aki\_stage:**

factor	AME	lower	upper	p
aki_stage2	1.011	-3.127	5.148	0.6322
aki_stage3	19.13	-1.16	39.43	0.06461

- persistent\_aki\_0\_transient\_aki\_1\_persistent\_aki:**

factor	AME	low er	up per	p
persistent_aki_0_transient_aki_1_persistent_aki1	7.201	0.5066	13.9	0.03501

- persistent\_aki\_stage:**

factor	AME	lower	upper	p
persistent_aki_stage1	2.889	-15.32	21.09	0.7558
persistent_aki_stage2	3	-23.57	29.57	0.8249
persistent_aki_stage3	19.33	-9.378	48.04	0.1869

- akd\_0\_no\_1\_yes:**

factor	AME	lower	upper	p
akd_0_no_1_yes1	6.302	-5.38	17.98	0.2903